arXiv:1411.2203v1 [math.NT] 9 Nov 2014

IF $p^a||n$ WHERE $n > 4$ IS THE ORDER OF A CIRCULANT HADAMARD MATRIX, THEN THE ORDER OF p MODULO n/p^a IS ODD

LUIS H. GALLARDO

Abstract. We proved recently (see [8]) the result on the title for odd prime divisors of such an n. The result implies for many $n's$, more precisely, for an infinity of n's with an arbitrary fixed number of prime divisors, the inexistence of circulant Hadamard matrices, and the inexistence of Barker sequences of length $n > 13$. The proof used a result of Arasu. It turns out that there is another, shorter proof, of the more general result that includes the prime $p = 2$. This new proof is based on a result of Brock (see [2, Theorem 3.1]), and besides that, requires just the definition of the Fourier transform. I noticed Brock's result in a preprint (see [12]) of Winterhof et al. where it is used to study the inexistence of related Butson-Hadamard matrices.

1. Introduction

A complex matrix H of order n is *complex Hadamard* if $HH^* = nI$, where I is the identity matrix of order n, and if every entry of $H/\sqrt{n}$ is in the complex unit circle. Here, the $*$ means transpose and conjugate. When such H has real entries, so that H is a $\{-1, 1\}$- matrix, H is called *Hadamard*. If H is Hadamard and circulant, say $H = circ(h_1, \ldots, h_n)$, that means that the i-th row H_i of H is given by $H_i = [h_{1-i+1}, \ldots, h_{n-i+1}]$, the subscripts being taken modulo n, for example $H_2 = [h_n, h_1, h_2, \ldots, h_{n-1}]$. A long standing conjecture of Ryser (see [11, pp. 134]) is:

Conjecture 1.1. *Let $n \geq 4$. If H is a circulant Hadamard matrix of order n, then $n = 4$.*

Details about previous results on the conjecture and a short sample of recent related papers are in [9], [10], [4], [3], [6], [7], [8].

The latter paper [8], based in [1], contains a proof of a practical criterion to reject many values of an integer n with $n > 4$, as possible orders of a circulant Hadamard matrix H.

The object of the present paper is to prove the following generalization of these result:

Theorem 1.2. *Let H be a circulant Hadamard matrix of order $n = p_1^{2a_1} \cdots p_t^{2a_t}$ where $p_1 = 2$, $a_1 = 1$ and $p_2, \ldots, p_t$ are all the distinct odd prime factors of n. Set for all $j = 1, \ldots, t$ $m_j := n/p_j^{2a_j}$. Then,*

$$o_m(p)$$

is odd, for any prime number p that divides n, where $m := m_j$ if $p := p_j$.

Date: November 11, 2014.

2000 *Mathematics Subject Classification.* Primary 11T55, 11T06; Secondary 11B73, 11B65, 05A10, 12E20.

Key words and phrases. Order of primes, Circulant Hadamard matrices.

The main result used in the proof is [2, Theorem 3.1] (see also Lemma 2.1 below). We discovered it in the recent preprint [12] of A. Winterhof, O. Yayla and V. Ziegler, where it is used to prove inexistence results on some Butson-Hadamard matrices (complex Hadamard matrices with all its entries k-th roots of 1 for some k).

2. Some tools

Brock's result (with some rewording) is the following.

Lemma 2.1. *Let $n = |a|^2 \in \mathbb{N}$ for some $a \in Q(w_{n_1})$, where $w_{n_1} = exp(2i\pi/n_1)$, then for every prime $p \nmid n_1$ such that $p \mid n$ the order $o_{n_1}(p)$ of p modulo n_1 is odd.*

Proof. See [2, Theorem 3.1]. □

The following is well known.

Lemma 2.2. *A circulant Hadamard matrix of order n exists if and only if $|\lambda| = \sqrt{n}$ for each eigenvalue λ of H.*

We recall the definition of the Fourier matrix, and in the lemma below the fundamental property of the Fourier transform for which we may check [5, pp .32–35].

Definition 2.3. Let n be a positive integer. Let $\omega := exp(2i\pi/n)$. Then, the Fourier matrix F is defined by the equality

$$\sqrt{n}F^* = Vandermonde([1, \omega, \omega^2, \ldots, \omega^{n-1}]) = (\omega^{(i-1)(j-1)}).$$

Lemma 2.4. *Let n be a positive integer. If $A = circ(a_1, \ldots, a_n)$ is a circulant matrix, of order n, with complex entries $a_j \in \mathbb{C}$ then its eigenvalues, b_j, $j = 1, \ldots, n$ in some order, are given by*

$$\sqrt{n} \cdot [\overline{a_1}, \ldots, \overline{a_n}] \cdot F = [\overline{b_1}, \ldots, \overline{b_n}]$$

where the $\bar{\cdot}$ denotes complex conjugation.

We are now ready to prove our main result in next section.

3. Proof of Theorem 1.2

Set $w_n := exp(2i\pi/n)$. Moreover, for any given $g \in \{1, \ldots, n\}$, put $n_1 := m_g$. Set $w_{n_1} := exp(2i\pi/n_1)$.

Let $H := circ(h_1, \ldots, h_n)$. Let $R(x) := h_1 + h_2x + \cdots + h_nx^{n-1}$ be the *representer* polynomial of H. By Lemma 2.4, the eigenvalues of H are the $b_s := R(w^{s-1})$ for all $s = 1, \ldots, n$. Define the index $j \in \{1, \ldots, n\}$ by

$$p_g^{2a_g} = j - 1. \tag{1}$$

Observe that

$$n = (2h)^2$$

where $h := 2 \cdot p_2^{a_2} \cdots p_t^{a_t}$. Now, by Lemma 2.2 one has also

$$n = b_j \cdot \overline{b_j} = |b_j|^2. \tag{2}$$

Observe that by definition of w_{n_1} and by (1) one has

$$w_{n_1} = w_n^{n/n_1} = w_n^{j-1}.$$

Thus,

$$b_j = R(w_n^{j-1}) = R(w_{n_1}) \in \mathbb{Q}(w_{n_1}). \tag{3}$$

Set $p := p_g$. Clearly $p \nmid n_1$ and $p \mid n$. Therefore, applying Lemma 2.1 to p one gets that

$$o_{n_1}(p)$$

is odd. This proves the theorem.

References

[1] K. T. Arasu, *A reduction theorem for circulant weighing matrices*, Australas. J. Combin. **18** (1998), p. 111–114.

[2] B. W. Brock, *Hermitian congruence and the existence and completion of generalized Hadamard matrices*, J. Combin. Theory Ser. A **49** (1988), no. 2, p. 233–261.

[3] R. A. Brualdi, *A note on multipliers of difference sets*, J. Res. Nat. Bur. Standards Sect. B **69** (1965), p. 87–89.

[4] R. Craigen, G. Faucher, R. Low, T. Wares, *Circulant partial Hadamard matrices*, Lin. Alg. Appl. **439**, 3307-3317, 2013.

[5] P. J. Davis, *Circulant matrices*, 2nd ed., New York, NY: AMS Chelsea Publishing, xix, 250 p. (1994).

[6] R. Euler, L. H. Gallardo, O. Rahavandrainy, *Sufficient conditions for a conjecture of Ryser about Hadamard Circulant matrices*, Lin. Alg. Appl. **437**, 2877-2886, 2012.

[7] L. Gallardo, *On a special case of a conjecture of Ryser about Hadamard circulant matrices*, Appl. Math. E-Notes **12**, 182-188, 2012.

[8] L. H. Gallardo, *Quadratic residues and a new infinity of orders for which a conjecture of Ryser about Circulant Hadamard matrices holds*, Preprint submitted, 2014.

[9] K. H. Leung, B. Schmidt, *New restrictions on possible orders of circulant Hadamard matrices*, Designs, Codes and Cryptography **64**, 143-151, 2012.

[10] M. J. Mossinghoff, *Wieferich pairs and Barker sequences*, Des. Codes Cryptogr. **53** (2009), 149–163.

[11] H. J. Ryser, Combinatorial mathematics. The Carus Mathematical Monographs, No. 14 Published by The Mathematical Association of America; distributed by John Wiley and Sons, Inc., New York 1963 xiv+154 pp.

[12] A. Winterhof, O. Yayla, V. Ziegler, *Non-existence of some nearly perfect sequences, near Butson-Hadamard matrices, and near Conference matrices*, Preprint, arXiv:1407.6548v1, 24Jul2014.

University Of Brest, Mathematics, 6, Av. Le Gorgeu, C.S. 93837, 29238 Brest Cedex 3, France.

E-mail address: Luis.Gallardo@univ-brest.fr